

+ 51% more closed wounds^{1*}

PICO[®] sNPWT[†] has been shown to significantly reduce wound area and depth when compared with tNPWT[‡] in patients with VLU and DFU over 12 weeks.¹

Turn around wound healing trajectory more effectively than standard dressings and tNPWT with PICO.²

Smith+Nephew



PICO[◇] 14

Single Use Negative Pressure Wound Therapy System

Helping you get **CLOSER TO ZERO[°]** delay in wound healing
smith-nephew.com/pico

References: 1. Kirsner R, Dove C, Reyzelman A, Vayser D, Jaimes H. A prospective, randomized, controlled clinical trial on the efficacy of a single-use negative pressure wound therapy system, compared to traditional negative pressure wound therapy in the treatment of chronic ulcers of the lower extremities. *Wound Rep Regen.* 2019. May 14 <https://doi.org/10.1111/wrr.12727>.
2. Dowsett C, et al. Use of PICO[®] to improve clinical and economic outcomes in hard-to-heal wounds. *Wounds International.* 2017;8, p53–58. * 45 vs 22%; p=0.002; ITT population. † Single Use Negative Pressure Wound Therapy (sNPWT). ‡ Traditional Negative Pressure Wound Therapy (tNPWT).
[°]Trademark of Smith & Nephew. All Trademarks acknowledged. ©October 2019 Smith & Nephew. AWM-AWD-20619 | CA48405 03/21